



University  
of Glasgow

## Bottom to Charm Decays with Heavy-HISQ

Judd Harrison, University of Glasgow

UKLFT Annual Meeting - May 2022



# Background


Many interesting  $B_{(s,c)}$  semileptonic decays with recent results/currently under active investigation

- ▶  $B_c \rightarrow D_s \ell^+ \ell^-$  and  $B_c \rightarrow D \ell \nu$ <sup>1</sup>
- ▶  $B_c \rightarrow D_s^* \ell^+ \ell^-$
- ▶  $B \rightarrow \pi, B \rightarrow K, B_s \rightarrow K$

---

<sup>1</sup>2111.06782

- Here, focus on three related  $b \rightarrow c$ , pseudoscalar to vector decays:  
 $B_{(s)} \rightarrow D_{(s)}^* \ell \nu$  and  $B_c \rightarrow J/\psi \ell \nu$
- General outline of the heavy-HISQ method, applied to form factor calculations,
  - Complementary determinations of  $V_{cb}$ ,
  - Comparison of observables sensitive to lepton flavor universality violation (LFUV) to experiment



Kinematic variables:

$$q^2 = (p - p')^2$$

$$w = \frac{p' \cdot p}{M_{B_q} M_{D_q^{(*)}}}$$

$$z = \frac{\sqrt{t_+ - q^2} - \sqrt{t_+ - t_0}}{\sqrt{t_+ - q^2} + \sqrt{t_+ - t_0}}$$

$$B_{(s)} \rightarrow D_{(s)}^* \ell \nu, B_c \rightarrow J/\psi \ell \nu$$

Pseudoscalar to vector decay has the following structure in the SM:

$$\frac{d\Gamma}{dq^2} = \chi(q^2) \times \mathcal{F}^2(q^2) |V_{cb}|^2$$

$$\mathcal{F}^2(q^2) = \left[ \left( 1 + \frac{m_\ell^2}{2q^2} \right) (H_+^2(q^2) + H_-^2(q^2) + H_0^2(q^2)) + \frac{3m_\ell^2}{2q^2} H_t^2(q^2) \right]$$

Helicity amplitudes expressed in terms of form factors

$$\{H_+(q^2), H_-(q^2), H_0(q^2)\} \leftrightarrow \{A_1(q^2), A_2(q^2), V(q^2)\}$$
$$H_t(q^2) \propto A_0(q^2)$$

- ▶ Theoretical predictions for vector meson final state require:
  - 4 form factors within the Standard Model
  - 3 additional tensor form factor for New Physics

- ▶  $V_{cb}$  - compare experimental value of  $\eta_{EW}\mathcal{F}(q_{\max}^2)|V_{cb}|$  to lattice calculations of  $\mathcal{F}(q_{\max}^2)$ 
  - preferred over  $B_{(s)} \rightarrow D_{(s)}$  due to favorable kinematics near zero-recoil.
  - Sensitive to choice of parameterisation scheme, preferable to compute using full kinematic range.
- ▶  $R(D^*) = \Gamma(B \rightarrow D^* \tau \bar{\nu}_\tau) / \Gamma(B \rightarrow D^* \mu \bar{\nu}_\mu)$ 
  - Sensitive to LFUV
  - Theory for  $R(D^*)$  relies on experimental fits + HQET for  $A_0$
  - On the lattice, typically use unphysically heavy pions and treat  $D^* \rightarrow D\pi$  resonance using  $\chi$ PT
- ▶ Lattice calculation of FFs for  $B_c \rightarrow J/\psi < B_s \rightarrow D_s^* < B \rightarrow D^*$ 
  - Computational cost of propagators for  $c < s \ll u/d$
  - $J/\psi$  and  $D_s^*$  are ‘gold-plated’
  - $B \rightarrow D^*$  requires careful treatment of chiral effects

# Overview of Lattice Results

- ▶ SM FFs for  $B \rightarrow D\ell\nu$  available away from zero recoil<sup>2</sup>
- ▶ SM FFs for  $B_s \rightarrow D_s\ell\nu$  now available across the full kinematic range, tensor FF available close to zero-recoil, with work also ongoing<sup>3</sup>
- ▶ SM FFs for  $B \rightarrow D^*\ell\nu$  recently became available from Fermilab-MILC away from zero-recoil<sup>4</sup>, with lattice calculations also underway by JLQCD as well as HPQCD.
- ▶ SM FFs for  $B_s \rightarrow D_s^*\ell\nu$  and  $B_c \rightarrow J/\psi\ell\nu$  available across full kinematic range from HPQCD<sup>5</sup>
- ▶ (Preliminary) SM FFs for  $B \rightarrow D^*\ell\nu$  across full kinematic range from HPQCD

---


<sup>2</sup>e.g. 1503.07237,1505.03925

<sup>3</sup>1906.00701,1310.5238,2110.10061


<sup>4</sup>2105.14019

<sup>5</sup>2105.11433, 2007.06957

# Current Results - $R(D^*)$



# Current Results - $V_{cb}$





## Current Results

	Lattice only	Lattice+Exp <sup>6</sup>	Experiment	Tension
$R(D)$	0.293(4) <sup>7</sup>	0.299(3)	0.340(30)	1.4 $\sigma$
$R(D^*)$	0.265(13)	0.2483(13)	0.295(14)	3.3 $\sigma$
$R(D_s)$	0.299(5)	—	—	—
$R(D_s^*)$	0.249(7)	—	—	—
$R(J/\psi)$	0.258(4)	—	0.71(25) <sup>8</sup>	1.8 $\sigma$

HFLAV average, Fermilab-MILC, HPQCD.


	$V_{cb}$	
$B \rightarrow D$	$39.58(94)_{\text{exp}}(37)_{\text{th}} \times 10^{-3}$	HFLAV
$B \rightarrow D^*$	$38.76(42)_{\text{exp}}(55)_{\text{th}} \times 10^{-3}$	
$B_s \rightarrow D_s^{(*)}$	$42.3(1.2)_{\text{exp}}(1.2)_{\text{th}} \times 10^{-3}$	LHCb (2001.03225)
$B \rightarrow X_c \ell \nu$	$42.16(51) \times 10^{-3}$	Bordone et al.(2107.00604)

<sup>6</sup>Assumes new physics only possible in semitauonic mode

<sup>7</sup>FLAG review

<sup>8</sup>LHCb-1711.05623

# Experimental Outlook



- ▶ Need precise SM form factors across full kinematic range
  - Resolve discrepancy between inclusive and exclusive determinations of  $V_{cb}$
  - Make first principles predictions for  $R(D_{(s)}^*)$  independent of experimental measurements
- ▶ Need tensor form factors to disentangle possible new physics effects

## $b \rightarrow c$ Pseudoscalar to Vector Form Factors

In the standard model  $\mathcal{F}(q^2)$  is a simple function of the form factors,  $A_1(q^2)$ ,  $A_0(q^2)$ ,  $A_2(q^2)$  and  $V(q^2)$ , defined in terms of matrix elements. For example, for  $B_s \rightarrow D_s^* \ell \nu$ :

$$\begin{aligned}\langle D_s^*(p', \lambda) | \bar{c} \gamma^\mu b | B_s^0(p) \rangle &= \frac{2iV(q^2)}{M_{B_s} + M_{D_s^*}} \epsilon^{\mu\nu\rho\sigma} \epsilon_\nu^*(p', \lambda) p'_\rho p_\sigma \\ \langle D_s^*(p', \lambda) | \bar{c} \gamma^\mu \gamma^5 b | B_s^0(p) \rangle &= 2M_{D_s^*} A_0(q^2) \frac{\epsilon^*(p', \lambda) \cdot q}{q^2} q^\mu \\ &+ (M_{B_s} + M_{D_s^*}) A_1(q^2) \left[ \epsilon^{*\mu}(p', \lambda) - \frac{\epsilon^*(p', \lambda) \cdot q}{q^2} q^\mu \right] \\ &- A_2(q^2) \frac{\epsilon^*(p', \lambda) \cdot q}{M_{B_s} + M_{D_s^*}} \left[ p^\mu + p'^\mu - \frac{M_{B_s}^2 - M_{D_s^*}^2}{q^2} q^\mu \right]\end{aligned}$$

# Form Factors Across the Full $q^2$ Range with Lattice QCD<sup>10</sup>

Use "Heavy-HISQ" approach:

- ▶ Compute form factors using multiple heavy masses ranging up to close to the physical b-quark mass
- ▶ Use **H**ighly **I**mproved **S**taggered **Q**uark action<sup>9</sup> for all quarks - fully relativistic, small discretisation effects
- ▶ Nonperturbatively renormalised currents, using PCVC and PCAC relations for vector and axial-vector, RI-SMOM for tensor
- ▶ Fit the form factor data including  $am_h$  discretisation effects, physical heavy mass dependence, and lattice spacing dependence
  - For  $B_s \rightarrow D_s^*$  and  $B_c \rightarrow J/\psi$  first convert to  $z$  space, e.g.

$$P(q^2) \times A_1(q^2) = \sum_{n=0}^3 a_n z^n(q^2) \mathcal{N}_n$$


$$a_n = \sum_{j,k,l=0}^3 b_n^{jkl} \left( \frac{2\Lambda_{\text{QCD}}}{M_{\eta_h}} \right)^j \left( \frac{am_c^{\text{val}}}{\pi} \right)^{2k} \left( \frac{am_h^{\text{val}}}{\pi} \right)^{2l}$$

---


<sup>9</sup>hep-lat/0610092

<sup>10</sup> $B_s \rightarrow D_s^*$ :2105.11433,  $B_c \rightarrow J/\psi$ :2007.06957

- ▶ We use the second generation MILC HISQ gauge configurations with  $u/d$ ,  $s$  and  $c$  quarks in the sea.





- ▶ The subset of configurations we use include physical  $u/d$  quark masses, and have small lattice spacings allowing us to come very close to the physical  $b$  mass.



$P(q^2) \times A_1$  for  $B_c \rightarrow J/\psi$ , plotted in  $z$  space, showing the physical continuum form factor as a blue band

# $B_c \rightarrow J/\psi$ Results - 2007.06956, 2007.06957




$$R(J/\psi) = 0.2582(38)$$

$$\Gamma(B_c^- \rightarrow J/\psi \mu^- \bar{\nu}_\mu) / \eta_{\text{EW}}^2 |V_{cb}|^2 = 1.73(12) \times 10^{13} \text{ s}^{-1}$$

- ▶ Experimental results for  $B_c \rightarrow J/\psi$  are currently much less precise than our lattice results, but expect this to improve in future.
- ▶ In addition to  $R(J/\psi)$ , other observables and ratios may be constructed with high precision from our form factor results
  - Can study the effect of NP couplings - full details in 2007.06956



# $B_s \rightarrow D_s^*$ Results - 2105.11433



$$R(D_s^*) = 0.249(6)_{\text{latt}}(4)_{\text{EM}}$$

$$\Gamma(B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu) / \eta_{\text{EW}}^2 |V_{cb}|^2 = 2.06(16) \times 10^{13} \text{s}^{-1}$$

$$R(D_s^*), V_{cb} \dots$$

Many new lattice predictions for  $B_s \rightarrow D_s^*$  quantities:

	This work	Exp. <sup>11</sup>	$B \rightarrow D^*$ <sup>12</sup>
$\frac{\Gamma(B_s^0 \rightarrow D_s^- \mu^+ \nu_\mu)}{\Gamma(B_s^0 \rightarrow D_s^{*-} \mu^+ \nu_\mu)}$	0.443(40)	0.464(45)	0.457(23)
$R(D_{(s)}^*)$	0.249(7)	—	0.2483(13)
$F_L$	0.440(16)	—	0.464(10)
$\mathcal{A}_{\lambda_\tau} = -P_\tau$	0.520(12)	—	0.496(15)

- ▶ Can also infer a total experimental rate  $\Gamma$  from LHCb analysis of  $V_{cb}$  in 2001.03225, we can use this with our results to give a value of  $V_{cb}$

$$|V_{cb}| = 42.2(2.3) \times 10^{-3}$$


- ▶ Consistent with the result using lattice data only at zero-recoil.

<sup>11</sup>LHCb 2001.03225

<sup>12</sup>HFLAV 1909.12524, Bordone et. al 1908.09398

## $B_s \rightarrow D_s^*$ Shape

We can compare the binned experimental differential rate<sup>13</sup> for the  $B_s \rightarrow D_s^*$  shape to our results




$$\chi^2/\text{dof} = 1.8 \text{ (0.62 excluding third bin)}$$

<sup>13</sup>LHCb:2003.08453

## $B_s \rightarrow D_s^*$ Shape Parameters

In the CLN parameterisation, the shape of the decay for massive leptons in the SM is fully described by the four parameters  $\rho^2$ ,  $R_1(1)$ ,  $R_2(1)$  and  $R_0(1)$ , with  $\rho^2$ ,  $R_1(1)$ ,  $R_2(1)$  determined from experiment and  $R_0(1)$  known to NLO in HQET<sup>14</sup>



- ▶ Our results are broadly consistent with the measured values of  $\rho^2$ ,  $R_1(1)$  and  $R_2(1)$  for  $B_s \rightarrow D_s^*$ , and with the NLO HQET value of  $R_0(1)$ .

<sup>14</sup>LHCb:2001.03225+2003.08453, HFLAV:1909.12524, HQET:1703.05330

## Preliminary results for $B \rightarrow D^*$

For  $B \rightarrow D^*$ , use HQET form factors:

$$\frac{\langle D^*(p', \lambda) | \bar{c} \gamma^\mu b | B(p) \rangle}{\sqrt{M_B M_{D^*}}} = h_V(w) \varepsilon^{\mu\nu}{}_{\rho\sigma} \epsilon_\nu^*(v_{D^*}, \lambda) v_{D^*}^\rho v_B^\sigma$$



$$\frac{\langle D^*(p', \lambda) | \bar{c} \gamma^\mu \gamma^5 b | B(p) \rangle}{\sqrt{M_B M_{D^*}}} = i \epsilon_\nu^* [g^{\mu\nu}(w+1) h_{A_1}(w) - v_B^\nu (v_B^\mu h_{A_2}(w) + v_{D^*}^\mu h_{A_3}(w))]$$

Computation completed on  $a = 0.045\text{fm}$ ,  $a = 0.06\text{fm}$  and  $a = 0.09\text{fm}$   $m_l = m_s/5$  lattices and  $a = 0.09\text{fm}$  physical  $m_l$  lattices. In the process of generating correlation functions on  $a = 0.06\text{fm}$  physical  $m_l$  lattices. Fit form factors to HQET inspired form, including chiral terms:

$$F = \sum_{nijk} a_{ijk}^n (w-1)^n \left(\frac{am_c}{\pi}\right)^i \left(\frac{am_h}{\pi}\right)^j \left(\frac{\Lambda_{\text{QCD}}}{M_B}\right)^k \mathcal{N}_n$$



$$+ X_{\log}(M_\pi/\Lambda_\chi) + A \left(\frac{M_\pi}{\Lambda_\chi}\right)^2$$

# Preliminary results for $B \rightarrow D^*$



We include data from  $B_s \rightarrow D_s^*$  in our chiral extrapolation.

# Preliminary results for $B \rightarrow D^*$




$$\left. \frac{d\mathcal{F}}{dw} \right|_{w=1} = -0.96(9)$$

$$R(D^*) = 0.276(13)$$

(PRELIMINARY)

(PRELIMINARY)

# Preliminary results for $B \rightarrow D^*$





Joint fit to HPQCD lattice and Belle untagged data -  $\chi^2/\text{dof} = 1.6$



$$V_{cb} = 39.3(0.7)_{\text{latt}}(0.5)_{\text{exp}} \times 10^{-3} \quad (\text{PRELIMINARY})$$



# Comparison to Fermilab-MILC (2105.14019)



# Comparison to Fermilab-MILC (2105.14019)



# Future Work

- ▶ Extend to unstable final states,  $B_s \rightarrow \phi$ ,  $B \rightarrow K^*$
- ▶ Extend to baryonic decays,  $\Lambda_b \rightarrow \Lambda_c$ 
  - 6 SM FFs, 4 additional tensor FFs
  - Complementary determination of  $V_{cb}$
  - $R(\Lambda_c)$

## Summary

- ▶ Published lattice results for  $B_c \rightarrow J/\psi$  form factors, corresponding experimental measurements are currently imprecise, but expected to improve.
- ▶ Results for the  $B_s \rightarrow D_s^*$  form factors on arXiv.
  - Model independent determinations of  $R(D_s^*)$  and other observables
  - Model independent determination of  $|V_{cb}|$ , though ideally would use experimental results directly
- ▶ Work on  $B \rightarrow D^*$  form factors almost complete, including tensor FFs and update to  $B_c \rightarrow J/\psi$  and  $B_s \rightarrow D_s^*$  to include tensor FFs
  - Preliminary value of  $V_{cb}$  consistent with existing exclusive determinations
  - Preliminary lattice only determination of  $R(D^*)$  higher than lattice+experiment

Thanks for listening!